

The synchrotron-self-Compton radiation accompanying shallow decaying X-ray afterglow: the case of GRB 940217

Da-Ming Wei^{1,2,3} and Yi-Zhong Fan^{1,2,4,5}

¹ *Purple Mountain Observatory, Chinese Academy of Science, Nanjing 210008, China.*

² *National Astronomical Observatories, Chinese Academy of Sciences, Beijing, 100012, China.*

³ *Joint Center for Particle Nuclear Physics and Cosmology of Purple Mountain Observatory - Nanjing University, Nanjing 210008, China.*

⁴ *The Racah Inst. of Physics, Hebrew University, Jerusalem 91904, Israel.*

⁵ *Golda Meir Fellow.*

ABSTRACT

High energy emission ($> \text{tens MeV}$) of Gamma-Ray Bursts (GRBs) provides an important clue to understand the physical processes involved in GRBs, which may be correlated with the GRB early afterglow. A shallow decline phase has been well detected in about half *Swift* Gamma-ray Burst X-ray afterglows. The widely considered interpretation involves a significant energy injection and possibly time-evolving shock parameter(s). This work we calculate the synchrotron-self-Compton (SSC) radiation of such an external forward shock and show that it could explain the well-known long term high energy (i.e., tens MeV to GeV) afterglow of GRB 940217. We propose that the cooperation of Swift and GLAST will help to reveal the nature of GRBs.

Subject headings: gamma-rays: bursts - ISM: jets and outflows - radiation mechanisms: nonthermal

1. Introduction

Among the high energy (above tens MeV) afterglow of Gamma-ray Burst (GRB) detected so far, that of GRB 940217 is the longest and also the most energetic one. The sub-GeV emission lasted more than 5000 seconds and it included also a 18 GeV photon (Hurley *et al.* 1994). The spectrum in the energy range 1 MeV to 18 GeV, cannot be fitted with a simple power law (see Fig. 3 of Hurley *et al.* 1994). A new spectral component in the energy range larger than several tens MeV is needed. This finding motivates many interesting ideas: (i) the interaction of ultra-relativistic protons with a dense cloud (Katz 1994),

(ii) synchrotron-self-Compton (SSC) scattering in early forward/reverse shocks or during the prompt emission (Mészáros & Rees 1994) and (iii) an electromagnetic cascade of TeV γ -rays in the infrared/microwave background (Plaga 1995). However, the two important observation facts: (a) the count rate of high energy photons is almost a constant (b) the typical energy of these photons is nearly unchanged, have not been satisfactorily reproduced.

To interpret the unusual high energy afterglow of GRB 940217, the physical processes involved in the early GRB afterglow phase is highly needed. The successful launch of the *Swift* satellite did open a new window to reveal what happens in the early GRB afterglow phase. As summarized in Zhang et al. (2006) and Nousek et al. (2006), in a canonical *Swift* GRB X-ray afterglow lightcurve some interesting features are emerging, including the very early sharp decline (i.e., phase-I), a shallow decline of the X-ray afterglow (i.e., phase-II), and the energetic X-ray flares (i.e., phase-V). The interpretation and the implication of these features have been discussed in great detail (see Mészáros 2006; Piran & Fan 2007; Zhang 2007 for recent reviews). For Phase-II, which interests us here, a widely considered explanation is a significant energy injection $dE_{\text{inj}}/dt \propto t^{-q}$ (see Zhang et al. 2006 and Nousek et al. 2006 and the references therein). However, the energy injection process, if there is, seems to be not all the story. As shown in Fan & Piran (2006), for some GRBs with good quality multi-wavelength afterglow data, the X-ray and optical light curves break chromatically and thus challenge the energy injection model (see also Panaitescu *et al.* 2006, Huang *et al.* . 2007). An assumption additional to the energy injection to solve such a puzzle is that the shock parameter(s) may be shock strength dependent (i.e., time-dependent).

In this work, we calculate the synchrotron-self-Compton (SSC) radiation of the external forward shock undergoing a significant energy injection. The shock parameters, ϵ_e and ϵ_B , the fraction of shock energy given to the shocked electrons and magnetic field, are assumed to be time-dependent. We show the high energy afterglow of GRB 940217 could be given rise to in such a scenario.

2. The SSC emission of the forward shock

In this section we calculate the synchrotron-self-Compton (SSC) radiation of the external forward shock undergoing a significant energy injection and evolving shock parameters.

2.1. The SSC emission in the standard fireball model

Synchrotron radiation. As usual, we assume that the electrons accelerated by the shock would follow the power law distribution $dn_e/d\gamma_e \propto \gamma_e^{-p}$ for $\gamma_e > \gamma_m$, where γ_m is the minimum Lorentz factor of shocked electrons (Sari et al. 1998). Then following Yost *et al.* (2003) and Fan & Piran (2006), the observed typical frequency of synchrotron radiation is $\nu_m = 4.2 \times 10^{14} (\frac{1+z}{2})^{1/2} \epsilon_{e,-1}^{1/2} \epsilon_{B,-2}^{1/2} E_{52}^{1/2} t_3^{-3/2}$ Hz. Where z is the redshift of the GRB, ϵ_e and ϵ_B are the energy fraction occupied by the electrons and magnetic field respectively, E is the isotropic energy, t is the observer time. Here the convention $Q_x = Q/10^x$ has been adopted in cgs units throughout the text, and we have taken the spectral index of the electron distribution $p = 2.5$. The observed cooling frequency is $\nu_c = 4.1 \times 10^{16} (\frac{1+z}{2})^{\frac{-1}{2}} \epsilon_{B,-2}^{-1/2} E_{52}^{-1/2} n^{-1} t_3^{-1/2} (1+Y)^{-2}$ Hz, where n is the surrounding medium density, $Y = [-1 + \sqrt{1 + 4x\epsilon_e/\epsilon_B}]/2$ is the Compton parameter, $x \simeq \min\{1, (\nu_m/\nu_c)^{(p-2)/2}\}$ (Sari & Esin 2001). The peak flux is $F_{\nu,max} = 3.2 (\frac{1+z}{2})^{1/2} \epsilon_{B,-2}^{1/2} E_{52} n^{1/2} D_{28}^{-2}$ mJy. Where D_{28} is the luminosity distance in units of 10^{28} cm. So in the fast cooling phase ($\nu_c < \nu_m$) the light curve is (Sari *et al.* 1998): $F_\nu \propto t^{1/6} (1+Y)^{2/3}$ for $\nu_c > \nu$, $F_\nu \propto t^{-1/4} (1+Y)^{-1}$ for $\nu_m > \nu > \nu_c$, and $F_\nu \propto t^{-(3p-2)/4} (1+Y)^{-1}$ for $\nu > \nu_m$. While in the slow cooling phase ($\nu_c > \nu_m$) the light curve is: $F_\nu \propto t^{1/2}$ for $\nu_m > \nu$, $F_\nu \propto t^{-3(p-1)/4}$ for $\nu_c > \nu > \nu_m$, and $F_\nu \propto t^{-(3p-2)/4} (1+Y)^{-1}$ for $\nu > \nu_c$.

Please note that in previous works the evolution of the Compton parameter Y has always been ignored. However, If $Y \gg 1$, then the effect of Y evolution should be considered. From the relation $x = \min\{1, (\nu_m/\nu_c)^{(p-2)/2}\}$ we know that, in the fast cooling phase, $x = 1$, then Y is independent of the time, while in the slow cooling phase Y should evolve with time. $x = (\nu_c/\nu_m)^{-(p-2)/2} \propto (1+Y)^{p-2} t^{-(p-2)/2}$, then we get

$$Y \simeq 10^{\frac{3-p}{4-p}} \left(\frac{1+z}{2}\right)^{\frac{p-2}{2(4-p)}} \epsilon_{e,-1}^{\frac{p-1}{4-p}} \epsilon_{B,-2}^{\frac{p-3}{4-p}} E_{52}^{\frac{p-2}{2(4-p)}} n^{\frac{p-2}{2(4-p)}} t_3^{-\frac{p-2}{2(4-p)}} \quad (1)$$

So we find $Y \propto t^{-(p-2)/2(4-p)}$, then $\nu_c \propto t^{(3p-8)/2(4-p)}$, and so we have $F_\nu \propto t^{\frac{3p^2-12p+4}{4(4-p)}}$ for $\nu > \nu_c$.

The SSC emission. The effect of SSC process on GRB afterglow emission has been discussed by several authors (e.g. Wei & Lu 1998, 2000; Sari & Esin 2001). The typical frequency of SSC emission is

$$\begin{aligned} \nu_m^{IC} &= 2\gamma_m^2 \nu_m = 3.36 \times 10^{21} \left(\frac{1+z}{2}\right)^{1/2} \\ &\quad \epsilon_{e,-1}^4 \epsilon_{B,-2}^{1/2} E_{52}^{3/4} n^{-1/4} t_3^{-9/4} \text{ Hz} \end{aligned} \quad (2)$$

Where $\gamma_m = \epsilon_e \left(\frac{p-2}{p-1}\right) \frac{m_p}{m_e} \Gamma \simeq 2 \times 10^3 \epsilon_{e,-1} E_{52}^{1/8} n^{-1/8} t_3^{-3/8}$ is the minimum Lorentz factor of shocked electrons, Γ is the bulk Lorentz factor, the cooling Lorentz factor $\gamma_c \simeq 2 \times 10^4 \epsilon_{B,-2}^{-1} E_{52}^{-3/8} n^{-5/8} \left(\frac{1+z}{2}\right)^{-1/2} t_3^{1/8} (1+Y)^{-1}$, then the cooling frequency of SSC emission is

$$\begin{aligned} \nu_c^{IC} &\approx 2\gamma_c^2 \nu_c = 3.28 \times 10^{25} \left(\frac{1+z}{2}\right)^{-3/2} \\ &\epsilon_{B,-2}^{-7/2} E_{52}^{-5/4} n^{-9/4} t_3^{-1/4} (1+Y)^{-4} \text{ Hz} \end{aligned} \quad (3)$$

The peak flux of the SSC emission is (Sari & Esin 2001)

$$\begin{aligned} F_m^{IC} &\simeq \frac{1}{3} n \sigma_T r F_{\nu, \max} = 1.2 \times 10^{-13} \left(\frac{1+z}{2}\right) \\ &\epsilon_{B,-2}^{1/2} E_{52}^{5/4} n^{5/4} t_3^{1/4} D_{28}^{-2} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ MeV}^{-1} \end{aligned} \quad (4)$$

Therefore, in the fast cooling phase, the light curve is

$$F_\nu \propto \begin{cases} t^{1/3} (1+Y)^{4/3}, & \nu_c^{IC} > \nu, \\ t^{1/8} (1+Y)^{-2}, & \nu_m^{IC} > \nu > \nu_c^{IC}, \\ t^{-(9p-10)/8} (1+Y)^{-2}, & \nu > \nu_m^{IC}, \end{cases} \quad (5)$$

and in the slow cooling phase, the light curve is

$$F_\nu \propto \begin{cases} t, & \nu_m^{IC} > \nu, \\ t^{-(9p-11)/8}, & \nu_c^{IC} > \nu > \nu_m^{IC}, \\ t^{-(9p-10)/8} (1+Y)^{-2}, & \nu > \nu_c^{IC}. \end{cases} \quad (6)$$

Again, if considering Y evolution, then, $F_\nu \propto t^{(9p^2-38p+24)/8(4-p)}$ for $\nu > \nu_c^{IC}$.

2.2. The SSC emission with energy injection and evolving shock parameters

A. The SSC emission with energy injection. In the standard fireball model, the shock energy E is assumed to be constant. However, there are increasing evidences that the shock energy may increase with time during some period. One good example is the discovery of the "shallow decay phase" in the early X-ray light curves of many GRBs, which is usually attributed to the energy injection (Zhang *et al.* 2006; Nousek *et al.* 2006). Now we consider the case that there is significant continuous energy injection into the fireball, so the fireball decelerates less rapidly and the afterglow emission will show a shallow decline. The dynamical evolution and the synchrotron radiation signature for energy injection have

been discussed by many authors (Rees & Mészáros 1998; Dai & Lu 1998; Cohen & Piran 1999; Zhang & Mészáros 2001a; Zhang *et al.* 2006; Nousek *et al.* 2006; Fan & Xu 2006).

Here we assume the energy injection takes a form $dE_{\text{inj}}/dt \propto t^{-q}$ (Cohen & Piran 1999; Zhang & Mészáros 2001), then the fireball energy evolves with time as $E \propto t^{1-q}$, the fireball radius $r \propto t^{(2-q)/4}$, the minimum electron Lorentz factor $\gamma_m \propto \Gamma \propto t^{-(2+q)/8}$, the observed typical frequency of synchrotron radiation $\nu_m \propto \gamma_m^2 B \Gamma \propto t^{-(2+q)/2}$, the cooling Lorentz factor $\gamma_c \propto (1+Y)^{-1} t^{(3q-2)/8}$, and the observed cooling frequency of synchrotron radiation $\nu_c \propto \gamma_c^2 B \Gamma \propto (1+Y)^{-2} t^{-(2-q)/2}$, the peak flux $F_{\nu, \text{max}} \propto N_e B \Gamma \propto t^{1-q}$. The typical frequency of SSC emission is $\nu_m^{IC} \propto \gamma_m^2 \nu_m \propto t^{-3(2+q)/4}$, the cooling frequency of SSC emission is $\nu_c^{IC} \propto \gamma_c^2 \nu_c \propto (1+Y)^{-4} t^{(5q-6)/4}$, and the peak flux of SSC radiation is $F_m^{IC} \simeq \frac{1}{3} n \sigma_T r F_{\nu, \text{max}} \propto t^{(6-5q)/4}$. Using these relations, we can get the light curves of synchrotron radiation and SSC emission with energy injection.

B. The SSC emission with evolving shock parameters. In the standard fireball model, the shock parameters ϵ_e and ϵ_B are assumed to be constant. However, it is also possible that these quantities may vary with time. Yost *et al.* (2003), Fan & Piran (2006) and Ioka *et al.* (2006) have discussed the afterglow emission with ϵ_e and ϵ_B being time-dependent. By modeling the afterglow of several GRBs, it was found that the values of ϵ_e and ϵ_B of the forward shock are quite different from that of reverse shock (Fan *et al.* 2002; Zhang, Kobayashi & Mészáros 2003; Kumar & Panaitescu 2003; Wei *et al.* 2006). We note that the forward shock is ultra-relativistic, while the reverse shock is mild-relativistic, so this result suggests that the shock parameters may be correlated with the strength of the shock.

Following Fan & Piran (2006), here we simply assume $\epsilon_e \propto \Gamma^{-a}$, $\epsilon_B \propto \Gamma^{-b}$, and since $\Gamma \propto t^{-(2+q)/8}$, so $\epsilon_e \propto t^{(2+q)a/8}$, $\epsilon_B \propto t^{(2+q)b/8}$. Then we can obtain the light curve of synchrotron radiation and SSC emission.

C. The SSC emission with both the above effects. It is also possible that during the shock evolution, both the shock energy and the shock parameters evolve with time. Based on the previous analysis, we can obtain the synchrotron radiation and SSC emission light curves easily. Table 1 gives the temporal index α of the afterglow emission, where $F_\nu \propto t^{-\alpha}$ is adopted. We define $\alpha = \alpha_0 + \alpha_E + \alpha_v + \alpha_Y$, where α_0 corresponds to the contribution of the standard emission, α_E represents the contribution of the energy injection, α_v stands for the contribution of evolving shock parameters, and α_Y comes from the evolution of Compton parameter Y . For example, if we only consider energy injection, then $\alpha = \alpha_0 + \alpha_E$. If only evolving shock parameters is considered, then $\alpha = \alpha_0 + \alpha_v$. If both the effects are considered, then $\alpha = \alpha_0 + \alpha_E + \alpha_v$. If $Y \gg 1$, then the term α_Y should be included.

3. The case of GRB940217

GRB940217 was a very famous burst for its long-lasting high energy afterglow emission (Hurley *et al.* 1994). The high energy photons ($E > 30$ MeV) were recorded for about 5400 seconds, including an 18 GeV photon ~ 4500 s after the low energy emission had ended. The 30 MeV to 30 GeV EGRET spectrum is $(1.3 \pm 0.4) \times 10^{-8} (E/86 \text{ MeV})^{-2.83 \pm 0.64} \text{ photons cm}^{-2} \text{ s}^{-1} \text{ MeV}^{-1}$, excluding the 18 GeV photon. By integrating this spectrum, the fluence at > 30 MeV is $7 \times 10^{-6} \text{ erg cm}^{-2}$. In addition, for this GRB there are two important observation facts: (a) the count rate of high energy photons is almost a constant; (b) the typical energy of these photons is nearly unchanged. This two facts imply that the flux should be nearly a constant $\sim 1.4 \times 10^{-9} \text{ erg cm}^{-2} \text{ s}^{-1}$.

Based on the above analysis, if we assume that the energy injection occurred at time $t \sim 500$ s, and we take the parameters as follows: $\epsilon_{e,-1} \sim 0.7$, $\epsilon_{B,-2} \sim 0.5$, $n \sim 1$, $E_{52} \sim 5$, $z \sim 0.1$, then at this time, the typical frequency of SSC emission is $\nu_m^{IC} \sim 20$ MeV, the cooling frequency of SSC emission is $\nu_c^{IC} \sim 60$ GeV, and the peak flux of SSC emission is $F_m^{IC} \sim 1.4 \times 10^{-11} \text{ ergs cm}^{-2} \text{ s}^{-1} \text{ MeV}^{-1}$, which is quite agreement with the observation. In addition, we note that the observed photon energies lie between ν_m^{IC} and ν_c^{IC} , and since the spectrum is very soft, $\beta = 1.83 \pm 0.64$, so the index of electron distribution is $p \sim 4$. From table 1 we can find that, in the standard case, the flux would decrease with time as $t^{-25/8}$, which is obviously inconsistent with the observation. If we consider energy injection, then $F_\nu \propto t^{-(6+19q)/8}$, since $0 < q < 1$, so the flux would decay steeper than $t^{-3/4}$, which is still inconsistent with the observation. Then, there is only one choice - the shock parameters should evolve with time. If we take $q \sim 0.5$ (which is suggested by the recent *Swift* XRT data, see Zhang *et al.* 2006), $a \sim 1$, $b \sim 0$, then the flux would be nearly a constant, which is agreement with the observation.

In order to investigate the SSC emission more carefully, Fan *et al.* (2007) have developed a numerical code to calculate the Compton process self-consistently for GRB high energy afterglow emission. Using this code, we calculate the SSC emission numerically, the result has been shown in Fig.1, the parameter are: the initial kinetic energy is 4×10^{52} erg, $n = 1$, $z = 0.1$, $\epsilon_{B,-2} = 0.3$, $p = 4$, and $\theta_j = 0.2$. For $10 \text{ s} < t < 5000 \text{ s}$, the energy injection form is taken to be $dE/dt = 4 \times 10^{50} (t/10 \text{ s})^{-0.55}$ and $\epsilon_e = 0.06 (t/5000 \text{ s})^{0.4}$. At late times, the energy injection disappears and $\epsilon_e = 0.06$. From Fig.1 we find that the numerical results are consistent with the analytic estimates and can well account for the observation (both the light curve and the spectrum) of GRB940217.

4. Discussion and conclusion

Since its discovery, the long-lasting high energy afterglow emission of GRB 940217 has been extensively discussed (e.g. Katz 1994; Mészáros & Rees 1994; Plaga 1995; Cheng & Cheng 1996; Dermer *et al.* 2000; Wang *et al.* 2001; Zhang & Mészáros 2001b; Guetta & Granot 2003; Pe’er & Waxman 2004). However, we note that the SSC emission of the standard afterglow model cannot well account for the observation of GRB940217. For example, Dermer *et al.* (2000) calculated the synchrotron-self-Compton emission during the blast wave propagation, but their spectrum is very hard, which is inconsistent with the observation of GRB 940217. As shown in Tab.1, the flux of SSC emission would decrease with time as $t^{-25/8}$ for a $p \sim 4$ that is needed to reproduce the very steep MeV to GeV spectrum, even when considering the energy injection, the flux would still decay steeper than $t^{-3/4}$. Therefore, the nearly constant of the high energy flux strongly suggests that other physical processes should be involved, such as the evolution of shock parameters and/or energy injection considered in this paper. Some possible energy injection processes have been proposed in Rees & Mészáros (1998) and Dai & Lu (1998), while the possible physical scenario giving rise to the time-evolving shock parameters are far from clear (Piran & Fan 2007). The peculiar chromatic break detected in quite a few early X-ray and optical afterglow data (Fan & Piran 2006; Panaitescu *et al.* 2006), however, did indicate such a possibility.

In previous analysis, the dependence of the Compton parameter Y on time has always been ignored. However, from table 1 we can see that, in some cases, the influence of Y evolution cannot be ignored. For example, in the standard case ($q = 1$, $a = b = 0$, i.e. without energy injection and the shock parameters are constant), the light curves of synchrotron radiation ($\nu > \nu_c$) and SSC emission ($\nu > \nu_c^{IC}$) will be flattened by $t^{1/6}$ and $t^{1/3}$ respectively for $p = 2.5$. If $p = 3$, then the effect will be more prominent, the light curves of synchrotron radiation ($\nu > \nu_c$) and SSC emission ($\nu > \nu_c^{IC}$) will be flattened by $t^{1/2}$ and t^1 respectively. So under some circumstances, the effect of Y evolution should be considered.

GLAST will be launched soon, and it is expected that GLAST will detect high energy emission (20 MeV to 300 GeV) of GRBs, which may open a new window to understand the physical processes occurred in GRBs. We hope that GLAST can detect more events like GRB940217, and this can provide important clues to explore the nature of GRBs (see Fan *et al.* 2007 for extensive discussions).

Acknowledgments

This work is supported by the National Natural Science Foundation (grants 10225314, 10233010, 10621303 and 10673034) of China.

REFERENCES

- Cheng, L. X. & Cheng, K. S., 1996, ApJ, 459, L79
- Cohen, E., & Piran, T., 1999, ApJ, 518, 346
- Dai, Z.G., & Lu, T., 1998, A&A, 333, L87
- Dermer, C.D., Chiang, J., & Mitman, K., 2000, ApJ, 537, 785
- Fan, Y.Z. *et al.* , 2007, MNRAS, to be submitted
- Fan, Y.Z., Dai, Z.G., Huang, Y.F., & Lu, T., 2002, Chin. J. Astron. Astrophys., 2, 449
- Fan, Y.Z., & Piran, T. 2006, MNRAS, 369, 197
- Fan, Y.Z., & Xu, D. 2006, MNRAS, 372, L19
- Guetta, D., & Granot, J., 2003, ApJ, 585, 885
- Huang, K. Y., *et al.*, ApJ, 654, L25
- Hurley K., *et al.* , 1994, Nature, 372, 652
- Ioka, K., Toma, K., Yamazaki, R., Nakamura, T., 2006, A&A, 458, 7
- Katz, J. I., 1994, ApJ, 432, L27
- Kumar, P., & Panaitescu, A., 2003, MNRAS, 346, 905
- Mészáros, P., 2006, Rep. Prog. Phys, 69, 2259
- Mészáros, P., & Rees, M. J., 1994, MNRAS, 269, L41
- Nousek, J. A. *et al.* 2006, ApJ, 642, 389
- Panaitescu, A. *et al.* , 2006, MNRAS, 369, 2059
- Pe’er, A., & Waxman, E., 2004, ApJ, 603, L1

- Plaga, R., 1995, *Nature*, 374, 430
- Piran, T., & Fan, Y. Z., 2007, *Phil. Trans. R. Soc. A.*, in press
- Rees, M. J., & Mészáros, P., 1998, *ApJ*, 496, L1
- Sari, R., & Mészáros, P., 2000, *ApJ*, 535, L33
- Sari, R., & Esin, A. A., 2001, *ApJ*, 548, 787
- Sari, R., Piran, T., & Narayan, R., 1998, *ApJ*, 497, L17
- Schaefer, B. E., *et al.*, 1998, *ApJ*, 492, 696
- Schneid, E. J., *et al.*, 1992, *A&A*, 255, L13
- Sommer, M., *et al.*, 1994, *ApJ*, 422, L63
- Wang, X. Y., Dai, Z. G., & Lu, T., 2001, *ApJ*, 556, 1010
- Wei, D. M., & Lu, T., 1998, *ApJ*, 505, 252
- Wei, D. M., & Lu, T., 2000, *A&A*, 360, L13
- Wei, D. M., Yan, T., Fan, Y. Z., 2006, *ApJ*, 636, L29
- Yost, S. A., Harrison, F. A., Sari, R., & Frail, D. A., 2003, *ApJ*, 597, 459
- Zhang, B., Mészáros, P., 2001a, *ApJ*, 552, L35
- Zhang, B., Mészáros, P., 2001b, *ApJ*, 559, 110
- Zhang, B., Kobayashi, S., & Mészáros, P., 2003, *ApJ*, 595, 950
- Zhang, B., Fan, Y. Z., Dyks, J., Kobayashi, S., Mészáros, P., *et al.* 2006, *ApJ*, 642, 354
- Zhang, B., 2007, *Chin. J. Astron. Astrophys.*, 7, 1

Table 1: The Temporal index α of afterglow emission. Here $F_\nu \propto t^{-\alpha}$ is adopted. We define $\alpha = \alpha_0 + \alpha_E + \alpha_v + \alpha_Y$, where α_0 corresponds to the contribution of standard emission, α_E represents the contribution of energy injection, α_v stands for the contribution of evolving shock parameters, and α_Y comes from the evolution of Compton parameter Y.

	α_0	α_E	α_v	α_Y
Synchrotron radiation	slow cooling			
$\nu < \nu_m$	$-\frac{1}{2}$	$-\frac{5(1-q)}{6}$	$\frac{(2a-b)(2+q)}{24}$	0
$\nu_m < \nu < \nu_c$	$\frac{3(p-1)}{4}$	$-\frac{(1-q)(p+3)}{4}$	$-\frac{4a(2+q)(p-1)+b(2+q)(p+1)}{32}$	0
$\nu > \nu_c$	$\frac{3p-2}{4}$	$-\frac{(1-q)(p+2)}{4}$	$-\frac{4a(2+q)(p-1)+b(2+q)(p-2)}{32}$	$-\frac{4q(p-2)-a(2+q)(p-1)-b(2+q)(p-2)}{8(4-p)}$
Synchrotron radiation	fast cooling			
$\nu < \nu_c$	$-\frac{1}{6}$	$-\frac{7(1-q)}{6}$	$-\frac{(2+q)b}{8}$	$-\frac{(2+q)(a-b)}{24}$
$\nu_c < \nu < \nu_m$	$\frac{1}{4}$	$-\frac{3(1-q)}{4}$	$\frac{(2+q)b}{32}$	$\frac{(2+q)(a-b)}{16}$
$\nu > \nu_m$	$\frac{3p-2}{4}$	$-\frac{(1-q)(p+2)}{4}$	$-\frac{4a(2+q)(p-1)+b(2+q)(p-2)}{32}$	$\frac{(2+q)(a-b)}{16}$
SSC emission	slow cooling			
$\nu < \nu_m^{IC}$	-1	$-(1-q)$	$\frac{(2+q)(4a-b)}{24}$	0
$\nu_m^{IC} < \nu < \nu_c^{IC}$	$\frac{9p-11}{8}$	$-\frac{(1-q)(3p+7)}{8}$	$-\frac{8a(2+q)(p-1)+b(2+q)(p+1)}{32}$	0
$\nu > \nu_c^{IC}$	$\frac{9p-10}{8}$	$-\frac{(1-q)(3p+2)}{8}$	$-\frac{8a(2+q)(p-1)-b(2+q)(6-p)}{32}$	$-\frac{4q(p-2)-a(2+q)(p-1)-b(2+q)(p-2)}{4(4-p)}$
SSC emission	fast cooling			
$\nu < \nu_c^{IC}$	$-\frac{1}{3}$	$-\frac{5(1-q)}{3}$	$-\frac{5b(2+q)}{24}$	$-\frac{(2+q)(a-b)}{12}$
$\nu_c^{IC} < \nu < \nu_m^{IC}$	$-\frac{1}{8}$	$-\frac{5(1-q)}{8}$	$\frac{5b(2+q)}{32}$	$\frac{(2+q)(a-b)}{8}$
$\nu > \nu_m^{IC}$	$\frac{9p-10}{8}$	$-\frac{(1-q)(3p+2)}{8}$	$-\frac{8a(2+q)(p-1)-b(2+q)(6-p)}{32}$	$\frac{(2+q)(a-b)}{8}$




Fig. 1.— The SSC radiation of the forward shock undergoing energy injection and with evolving shock parameters, the case of GRB 940217: the thick solid line is the light curve and the inserted plot is the spectrum (the times have been marked in the plot), in the energy range of 30 MeV - 30 GeV. For $10 \text{ s} < t < 5000 \text{ s}$, the energy injection form is taken to be $dE/dt = 4 \times 10^{50}(t/10\text{s})^{-0.55}$ and $\epsilon_e = 0.06(t/5000\text{s})^{0.4}$. At late times, the energy injection disappears and $\epsilon_e = 0.06$. Other parameters involved in the calculation are as follows: the initial kinetic energy is $4 \times 10^{52} \text{ erg}$, $z = 0.1$, $\epsilon_{B,-2} = 0.3$, $p = 4$, and $\theta_j = 0.2$.